

Spectroscopic Survey of Runaway Star Candidates


Aakash Bhat & Stephan Geier

What is a star?



Dmitry Brant/Wikimedia Commons

Gas balls in equilibrium



Gravitational force acting on shell
 r with thickness dr inward


$$G = -g\rho dr$$

Balanced by force due to pressure
outward (**Fusion processes**)

$$P_i - P_e = -\frac{\partial P}{\partial r} dr$$

Hydrostatic equilibrium


Hot Main Sequence Stars



Kippenhahn, Weigert & Weiss 2012


Spectral-type B

$T_{\text{eff}} = 10000 - 30000 \text{ K}$



X-Shooter spectral library, <http://xsl.u-strasbg.fr/>


Hot Main Sequence Stars




Kippenhahn, Weigert & Weiss 2012

Spectral-type O

$T_{\text{eff}} = 30000 - 50000 \text{ K}$



Hot Main Sequence Stars




The **nuclear lifetime** on the main sequence is a strong function of L and therefore M

$$\tau_H \sim M^{-2.5}$$

It ranges from several million years to more than the age of the Universe for $M < 0.8 M_\odot$


Hot MS stars are short-lived

Hot Main Sequence Stars




**Hot MS stars are
very rare**

Single low-mass stellar evolution




Thomas 1967, ZA, 67, 420

$1.3 M_{\odot}$ Radiative core (Low mass)




Degenerate helium core grows in mass due to central H-burning

Single low-mass stellar evolution



- In low-mass stars the core is radiative**
 - No efficient mixing in the core
 - Hydrogen is consumed starting in the center
 - Smooth transition to shell burning

Single low-mass stellar evolution




Due to the high density in the core,
the electron gas becomes **degenerate**

→ Isothermal, degenerate core is
stable

→ Core can grow in mass


Single low-mass stellar evolution



**No heating during core contraction
due to equation of state**


$$P_e = 1.0036 \times 10^{13} \left(\frac{\rho}{\mu_e} \right)^{5/3}$$

Single low-mass stellar evolution




Thomas 1967, ZA, 67, 420

$1.3 M_{\odot}$ Radiative core (**Low mass**)



H-shell burning starts → Core contracts, envelope expands


Single low-mass stellar evolution



Temperature of the core increases

- Increase of temperature in the H-burning shell
- Core contraction heats transition layer between core and shell


Single low-mass stellar evolution



Critical temperature for helium burning ($\sim 10^8$ K) is reached for a core mass of about $0.48 M_{\odot}$

Due to energy losses via neutrinos in the center, helium is ignited in a shell


Single low-mass stellar evolution



Due to the **high temperature dependency** of the 3α reaction rate $\langle\sigma v\rangle \sim \rho T^{40}$, nuclear energy is released fast and increases the core temperature


Degenerate gas cannot expand with increasing temperature

Single low-mass stellar evolution



Runaway burning of helium
Helium flash


Single low-mass stellar evolution



Runaway burning of helium under degenerate conditions

- Degeneracy is lifted
- Core expands, density drops
- Stable He-core burning


Single low-mass stellar evolution



Luminosity of the core
during the flash higher than
the luminosity of the Galaxy
($10^{11} L_\odot$)

→ Trapped in the
envelope

Horizontal branch stars




Phase of stable He-core and H-shell burning

→ Stars occupy a region of (about) constant luminosity

Horizontal Branch


Horizontal branch stars



Horizontal Branch stars

- Different mass loss η on the RGB leads to **different thickness of the hydrogen envelopes**
- Mass of the He-core is constant ($\sim 0.48 M_\odot$)
- Diverse types of HB stars


Horizontal branch stars



Horizontal Branch stars


- The thinner the hydrogen envelope, the bluer the HB star
- Morphology of HB depends on metallicity and age

Horizontal branch stars




Red Clump stars

- Red giants
- Intermediate mass stars
- Young population




Horizontal branch stars



Red Horizontal Branch (RHB) stars

- Redward of the MS
- (Sub-)giants
- Spectral types K, G
- metal-poor, old population


Horizontal branch stars



RR Lyr stars

- (Sub-)giants
- Spectral types F
- metal-poor, old population
- Pulsators


Horizontal branch stars



Blue Horizontal Branch (BHB) stars

- Blueward of the MS
- (Sub-)dwarfs
- Spectral types A, B (HBA, HBB)
- chemically peculiar

Horizontal branch stars




**Extreme Horizontal Branch (EHB)
stars**

→ Subdwarfs

→ Spectral types O, B (sdO, sdB)

→ Extremely thin hydrogen
envelopes, no H-shell burning

Horizontal branch stars




Hydrogen-rich sdBs

→ very low to solar helium content

→ Light elements depleted, heavy elements enriched

→ High binary fraction

Binary evolution




Binary fraction on the main sequence depends on stellar mass

~10 % triple

~1 % quadruple or higher multiple systems

Binary evolution



Raghavan et al. 2010, ApJS, 190, 1


Binary evolution



Stable mass transfer

**Common envelope
phase**


Binary evolution



Close binary evolution

- Helium-burning core of a red giant stripped by binary interaction
- Stable and unstable mass-transfer possible
- sdO/Bs predicted to be in close and wide binaries

Binary evolution




~30% of the sdO/Bs are in
**composite double-lined
binaries**

Companions are K/G/F-type
main sequence stars

The orbital periods of the
~30 solved systems
($P = 300 - 1200$ d) are in
the appropriate range for
prior RLOF mass-transfer

Binary evolution




Kupfer et al. 2015, A&A, 576, 44

~30% of the sdO/Bs are in single-lined close binaries

Companions are M-type main sequence stars, brown dwarfs and white dwarfs

The orbital periods of the ~300 solved systems ($P = 0.03 - 30$ d) are typical for post-CE systems

Binary evolution




ESA/ATG medialab

~30% of the sdO/Bs don't show any signs of binarity

- Close substellar companions such as brown dwarfs or planets
- Evaporation or merger during CE evolution?


Binary evolution



Helium-rich sdO/Bs


- very high helium abundance
- Enrichment in carbon and/or nitrogen
- Single stars

Binary evolution



NASA


Binary evolution



Alternative formation

- **Close binary evolution**
- **Merger of two white dwarfs of pure helium composition**
- **Single He-sdO/B stars**

Spectroscopic binaries



Spectral lines are shifted w.r.t.
their rest wavelengths

→ **Doppler effect**


$$\frac{\lambda - \lambda_0}{\lambda_0} = \frac{\Delta \lambda}{\lambda_0} = \frac{v}{c} \quad \text{for } v \ll c$$

λ observed wavelength

λ_0 rest wavelength


v radial velocity

Spectroscopic binaries



ESO


Spectroscopic binaries




Measuring line-shift

→ Radial velocity

Spectroscopic binaries



Spectroscopic binaries




RVs and times must be corrected for Earth's motion around the barycenter of the solar system (up to ± 30 km s^{-1} in RV and ± 8 min in time)

→ **Location of the telescope must be known (GPS)**

→ **Most accurate determination of observation time: High-speed photometers measure photon weighted midpoint of exposures**


Spectroscopic binaries



RVs and times must be corrected for Earth's motion around the barycenter of the solar system (up to ± 30 km s^{-1} in RV and ± 8 min in time)

- For close binaries with high RV shifts often slightly less accurate **heliocentric corrections** are used
- Times are approximated by adding **half of the exposure time** to the starting time


Spectroscopic binaries



Preparatory study at the workshop in 2021 and BHB survey in workshop 2022

→ RV accuracy of a few km/s

Spectroscopic binaries



Raghavan et al. 2010, ApJS, 190, 1

Preparatory study at the workshop in 2021 and BHB survey in workshop 2022

→ RV accuracy of a few km/s

→ Sensitive to orbital periods of several tens of days